

EXTENDED q -DEDEKIND-TYPE DAEHEE-CHANGHEE SUMS ASSOCIATED WITH EXTENDED q -EULER POLYNOMIALS

SERKAN ARACI AND MEHMET ACIKGOZ

ABSTRACT. In the present paper, our goal is to introduce a p -adic continuous function for an odd prime to inside a p -adic q -analogue of the higher order Dedekind-type sums with weight α related to Extended q -Euler polynomials by using p -adic q -integral in the p -adic integer ring.

2010 MATHEMATICS SUBJECT CLASSIFICATION. 11S80, 11B68.

KEYWORDS AND PHRASES. Dedekind Sums, q -Dedekind-type Sums, p -adic q -integral, Extended q -Euler numbers and polynomials.

1. Introduction

Assume that p be a fixed odd prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex numbers and the completion of algebraic closure of $\mathbb{Q}_p$.

Let v_p be normalized exponential valuation of $\mathbb{C}_p$ by

$$|p|_p = p^{-v_p(p)} = \frac{1}{p}.$$

When one talks of q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$ or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we assume that $|1 - q|_p < 1$ (see, [1-15]).

A q -analogue of p -adic Haar distribution is defined by Kim as follows: for any positive integer n ,

$$\mu_q(a + p^n \mathbb{Z}_p) = (-q)^a \frac{(1 + q)}{1 + q^{p^n}}$$

for $0 \leq a < p^n$ and this can be extended to a measure on $\mathbb{Z}_p$ (for details, see [1-7]).

Extended q -Euler polynomials are defined as

$$(1) \quad \tilde{E}_{n,q}^{(\alpha)}(x) = \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha(x+\xi)}}{1 - q^\alpha} \right)^n d\mu_q(\xi)$$

for $n \in \mathbb{Z}_+ := \{0, 1, 2, 3, \dots\}$. We note that

$$\lim_{q \rightarrow 1} \tilde{E}_{n,q}^{(\alpha)}(x) = E_n(x)$$

where $E_n(x)$ are n -th Euler polynomials, which are defined by the rule:

$$\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} = e^{tx} \frac{2}{e^t + 1}, \quad |t| < \pi$$

(for details, see [12]). Taking $x = 0$ into (1), then we have $\tilde{E}_{n,q}^{(\alpha)}(0) := \tilde{E}_{n,q}^{(\alpha)}$ are called extended q -Euler numbers.

Extended q -Euler numbers and polynomials have the following equalities

$$(2) \quad \tilde{E}_{n,q}^{(\alpha)} = \frac{1+q}{(1-q^\alpha)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^{\alpha l+1}},$$

$$(3) \quad \tilde{E}_{n,q}^{(\alpha)}(x) = \frac{1+q}{(1-q^\alpha)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{q^{\alpha l x}}{1+q^{\alpha l+1}},$$

$$(4) \quad \tilde{E}_{n,q}^{(\alpha)}(x) = \sum_{l=0}^n \binom{n}{l} q^{\alpha l x} \tilde{E}_{l,q}^{(\alpha)} \left(\frac{1-q^{\alpha x}}{1-q^\alpha} \right)^{n-l}.$$

Also, for $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$

$$(5) \quad \tilde{E}_{n,q}^{(\alpha)}(x) = \left(\frac{1+q}{1+q^d} \right) \left(\frac{1-q^{\alpha d}}{1-q^\alpha} \right)^n \sum_{a=0}^{d-1} (-1)^a \tilde{E}_{n,q}^{(\alpha)} \left(\frac{x+a}{d} \right),$$

(for more information, see [12]).

For any positive integer h, k and m , Dedekind-type DC sums are given by Kim in [1], [2] and [3] as follows:

$$S_m(h, k) = \sum_{M=1}^{k-1} (-1)^{M-1} \frac{M}{k} \overline{E}_m \left(\frac{hM}{k} \right)$$

where $\overline{E}_m(x)$ are the m -th periodic Euler function.

Recently, weighted q -Bernoulli numbers and polynomials was firstly defined by Taekyun Kim in [8]. After, many mathematicians, by utilizing from Kim's paper [8], have introduced a new concept in Analytic numbers theory as weighted q -Bernoulli, weighted q -Euler, weighted q -Genocchi polynomials in [11], [12], [14] and [15]. Also, Kim derived some interesting properties for Dedekind-type DC sums. He considered a p -adic continuous function for an odd prime number to contain a p -adic q -analogue of the higher order Dedekind-type DC sums $k^m S_{m+1}(h, k)$ in [2]. In [9], Simsek also studied to q -analogue of Dedekind-type sums. He also derived their interesting properties.

By the same motivation, we, by using p -adic q -integral on $\mathbb{Z}_p$, will construct weighted p -adic q -analogue of the higher order Dedekind-type DC sums $k^m S_{m+1}(h, k)$.

2. Extended q -Dedekind-type Sums associated with Extended q -Euler polynomials

Let w be the *Teichmüller* character $(\text{mod } p)$. For $x \in \mathbb{Z}_p^* := \mathbb{Z}_p/p\mathbb{Z}_p$, set

$$\langle x : q \rangle = w^{-1}(x) \left(\frac{1-q^x}{1-q} \right).$$

Let a and N be positive integers with $(p, a) = 1$ and $p \mid N$. We now consider the following

$$\tilde{C}_q^{(\alpha)}(s, a, N : q^N) = w^{-1}(a) \langle x : q^\alpha \rangle^s \sum_{j=0}^{\infty} \binom{s}{j} q^{\alpha a j} \left(\frac{1-q^{\alpha N}}{1-q^{\alpha a}} \right)^j \tilde{E}_{j,q^N}^{(\alpha)}.$$

In particular, if $m + 1 \equiv 0 \pmod{p - 1}$, then

$$\begin{aligned} \tilde{C}_q^{(\alpha)}(m, a, N : q^N) &= \left(\frac{1 - q^{\alpha a}}{1 - q^\alpha} \right)^m \sum_{j=0}^m \binom{m}{j} q^{\alpha a j} \tilde{E}_{j, q^N}^{(\alpha)} \left(\frac{1 - q^{\alpha N}}{1 - q^{\alpha a}} \right)^j \\ &= \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(\xi + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m d\mu_{q^N}(\xi). \end{aligned}$$

Thus, $\tilde{C}_q^{(\alpha)}(m, a, N : q^N)$ is a continuous p -adic extension of

$$\left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \tilde{E}_{m, q^N}^{(\alpha)} \left(\frac{a}{N} \right).$$

Let $[\cdot]$ be the Gauss' symbol and let $\{x\} = x - [x]$. Thus, we are now ready to introduce q -analogue of the higher order Dedekind-type DC sums $\tilde{J}_{m, q}^{(\alpha)}(h, k : q^l)$ by the rule:

$$\tilde{J}_{m, q}^{(\alpha)}(h, k : q^l) = \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha k}} \right) \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha(l\xi + l\{\frac{hM}{k}\})}}{1 - q^{\alpha l}} \right)^m d\mu_{q^l}(\xi).$$

If $m + 1 \equiv 0 \pmod{p - 1}$

$$\begin{aligned} &\left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha k}} \right) \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(\xi + \frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m d\mu_{q^k}(\xi) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^\alpha} \right) \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(\xi + \frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m d\mu_{q^k}(\xi) \end{aligned}$$

where $p \mid k$, $(hM, p) = 1$ for each M . Thanks to (1), we easily state the following

$$\begin{aligned} (6) \quad &\left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \tilde{J}_{m, q}^{(\alpha)}(h, k : q^k) \\ &= \sum_{M=1}^{k-1} \left(\frac{1 - q^{\alpha M}}{1 - q^\alpha} \right) \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^m (-1)^{M-1} \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(\xi + \frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m d\mu_{q^k}(\xi) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^\alpha} \right) \tilde{C}_q^{(\alpha)}(m, (hM)_k : q^k) \end{aligned}$$

where $(hM)_k$ denotes the integer x such that $0 \leq x < n$ and $x \equiv \alpha \pmod{k}$.

It is not difficult to indicate the following

$$\begin{aligned} (7) \quad &\int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha(x+\xi)}}{1 - q^\alpha} \right)^k d\mu_q(\xi) \\ &= \left(\frac{1 - q^{\alpha m}}{1 - q^\alpha} \right)^k \frac{1 + q}{1 + q^m} \sum_{i=0}^{m-1} (-1)^i \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha m(\xi + \frac{x+i}{m})}}{1 - q^{\alpha m}} \right)^k d\mu_{q^m}(\xi). \end{aligned}$$

On account of (6) and (7), we easily see that

$$(8) \quad \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(\xi + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m d\mu_{q^N}(\xi) \\ = \frac{1 + q^N}{1 + q^{Np}} \sum_{i=0}^{p-1} (-1)^i \left(\frac{1 - q^{\alpha Np}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha p N(\xi + \frac{a+iN}{pN})}}{1 - q^{\alpha p N}} \right)^m d\mu_{q^{pN}}(\xi).$$

Because of (6), (7) and (8), we develop the p -adic integration as follows:

$$\tilde{C}_q^{(\alpha)}(s, a, N : q^N) = \frac{1 + q^N}{1 + q^{Np}} \sum_{\substack{0 \leq i \leq p-1 \\ a+iN \not\equiv 0 \pmod{p}}} (-1)^i \tilde{C}_q^{(\alpha)}\left(s, (a + iN)_{pN}, p^N : q^{pN}\right).$$

So,

$$\tilde{C}_q^{(\alpha)}(m, a, N : q^N) = \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(\xi + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m d\mu_{q^N}(\xi) \\ - \left(\frac{1 - q^{\alpha Np}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha p N(\xi + \frac{a+iN}{pN})}}{1 - q^{\alpha p N}} \right)^m d\mu_{q^{pN}}(\xi)$$

where $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$ and m is integer with $m+1 \equiv 0 \pmod{p-1}$. Therefore, we procure the following

$$\sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^\alpha} \right) \tilde{C}_q^{(\alpha)}(m, hM, k : q^k) \\ = \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \tilde{J}_{m,q}^{(\alpha)}(h, k : q^k) - \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \left(\frac{1 - q^{\alpha kp}}{1 - q^{\alpha k}} \right) \tilde{J}_{m,q}^{(\alpha)}((p^{-1}h), k : q^{pk})$$

where $p \nmid k$ and $p \nmid hm$ for each M . Thus, we give the following definition, which seems interesting for further studying in theory of Dedekind sums.

Definition 1. Let h, k be positive integer with $(h, k) = 1$, $p \nmid k$. For $s \in \mathbb{Z}_p$, we define p -adic Dedekind-type DC sums as follows:

$$\tilde{J}_{p,q}^{(\alpha)}(s : h, k : q^k) = \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^\alpha} \right) \tilde{C}_q^{(\alpha)}(m, hM, k : q^k).$$

As a result of the above definition, we state the following theorem.

Theorem 2.1. For $m+1 \equiv 0 \pmod{p-1}$ and $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$, then, we have

$$\tilde{J}_{p,q}^{(\alpha)}(s : h, k : q^k) = \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \tilde{J}_{m,q}^{(\alpha)}(h, k : q^k) \\ - \left(\frac{1 - q^{\alpha k}}{1 - q^\alpha} \right)^{m+1} \left(\frac{1 - q^{\alpha kp}}{1 - q^{\alpha k}} \right) \tilde{J}_{m,q}^{(\alpha)}((p^{-1}h), k : q^{pk}).$$

In the special case $\alpha = 1$, our applications in theory of Dedekind sums resemble Kim's results in [2]. These results seem to be interesting for further studies in [1], [3] and [9].

REFERENCES

- [1] T. Kim, A note on p -adic q -Dedekind sums, *C. R. Acad. Bulgare Sci.* 54 (2001), 37–42.
- [2] T. Kim, Note on q -Dedekind-type sums related to q -Euler polynomials, *Glasgow Math. J.* 54 (2012), 121–125.
- [3] T. Kim, Note on Dedekind type DC sums, *Adv. Stud. Contemp. Math.* 18 (2009), 249–260.
- [4] T. Kim, The modified q -Euler numbers and polynomials, *Adv. Stud. Contemp. Math.* 16 (2008), 161–170.
- [5] T. Kim, q -Volkenborn integration, *Russ. J. Math. Phys.* 9 (2002), 288–299.
- [6] T. Kim, On p -adic interpolating function for q -Euler numbers and its derivatives, *J. Math. Anal. Appl.* 339 (2008), 598–608.
- [7] T. Kim, On a q -analogue of the p -adic log gamma functions and related integrals, *J. Number Theory* 76 (1999), 320–329.
- [8] T. Kim, On the weighted q -Bernoulli numbers and polynomials, *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 207–215, 2011.
- [9] Y. Simsek, q -Dedekind type sums related to q -zeta function and basic L -series, *J. Math. Anal. Appl.* 318 (2006), 333–351.
- [10] M. Acikgoz, Y. Simsek, On multiple interpolation function of the Nörlund-type q -Euler polynomials, *Abst. Appl. Anal.* 2009 (2009), Article ID 382574, 14 pages.
- [11] S. H. Rim and J. Jeong, A note on the Modified q -Euler Numbers and polynomials with weight α , *International Mathematical Forum*, Vol. 6, 2011, no. 65, 3245–3250.
- [12] C. S. Ryoo, A note on the weighted q -Euler numbers and polynomials, *Adv. Stud. Contemp. Math.* 21 (2011), page 47–54
- [13] S. Araci, M. Acikgoz, K. H. Park and H. Jolany, On the unification of two families of multiple twisted type polynomials by using p -adic q -integral on $\mathbb{Z}_p$ at $q = -1$, accepted in *Bulletin of the Malaysian Mathematical Sciences and Society*.
- [14] S. Araci, M. Acikgoz and K. H. Park, A note on the q -analogue of Kim's p -adic log gamma type functions associated with q -extension of Genocchi and Euler numbers with weight α , accepted in *Bulletin of the Korean Mathematical Society*.
- [15] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p -adic q -integral representation on $\mathbb{Z}_p$ associated with weighted q -Bernstein and q -Genocchi polynomials, *Abstract and Applied Analysis*, Volume 2011, Article ID 649248, 10 pages.
- [16] S. Araci, M. Acikgoz and J. J. Seo, Explicit formulas involving q -Euler numbers and polynomials, *Abstract and Applied Analysis*, Volume 2012, Article ID 298531, 11 pages (Article in press).

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY

E-mail address: mtsrkn@hotmail.com; saraci88@yahoo.com.tr; mtsrkn@gmail.com

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY

E-mail address: acikgoz@gantep.edu.tr